

## References

1. Lin RY, Shah SN. Increasing hospitalizations due to angioedema in the United States. *Ann Allergy Asthma Immunol* 2008;101:185-192.
2. Byrd JB, Adam A, Brown NJ. Angiotensin-converting enzyme inhibitor-associated angioedema. *Immunol Allergy Clin N Am* 2006;26:725-737.
3. Lo KS. Angioedema associated with candesartan. *Pharmacotherapy* 2002;22(9):1176-1179.
4. Orr KK, Myers JR. Intermittent visceral edema induced by long-term enalapril administration. *Ann Pharmacother* 2004;38(5):825-827.
5. Irons BK, Kumar A. Valsartan-induced angioedema. *Ann Pharmacother* 2003;37:1024-1027.
6. Schmidt TD, McGrath KM. Angiotensin-converting enzyme inhibitor angioedema of the intestine: a case report and review of the literature. *Am J Med Sci* 2002;324(2):106-108.
7. Agostoni A, Cicardi M. Drug-induced angioedema without urticaria. *Drug Saf* 2001;24(8):599-606.
8. Abdi R, Dong VM, Lee CJ, et al. Angiotensin II receptor blocker-associated angioedema: on the heels of ACE inhibitor angioedema. *Pharmacotherapy* 2002;22(9):1173-1175.
9. Greaves M, Lawlor F. Angioedema: manifestations and management. *J Am Acad Dermatol* 1991;25:155-165.
10. Cicardi M, Zingale LC, Bergamaschini L, et al. Angioedema associated with angiotensin-converting enzyme inhibitor use: outcome after switching to a different treatment. *Arch Intern Med* 2004;164:910-913.
11. Boehringer SK. Can angiotensin receptor blockers be used in patients who have previous ACE inhibitor-induced angioedema? *Pharmacist's Letter Detail-Document* 2004;#200709.
12. Pfeffer MA, McMurray JJ, Velazquez EJ, et al. Valsartan, captopril, or both in myocardial infarction complicated by heart failure, left ventricular dysfunction, or both. *N Engl J Med* 2003;349:1893-1906.